

Low Birth Weight Rate (<2,500 Grams, Excluding <500 Grams)

| | |
|--------------------------------|---|
| Name | Low Birth Weight Rate (<2,500 Grams, Excluding <500 Grams) |
| Short/Other Names | Not applicable |
| Description | Crude low birth weight rate (<2,500 grams, excluding <500 grams) |
| Interpretation | Lower rates for this indicator are desirable. |
| HSP Framework Dimension | Health System Outcomes: Improve health status of Canadians |
| Areas of Need | Not applicable |
| Geographic Coverage | All provinces/territories |
| Reporting Level/Disaggregation | National, Province/Territory, Region |
| Indicator Results | https://apps.cihi.ca/mstrapp/asp/Main.aspx?Server=apmstrextpd_i&project=Quick%20Stats&uid=pce_pub_en&pwd=&evt=2048001&visualizationMode=0&documentID=029DB170438205AEBCC75B8673CCE822 |

Identifying Information

Name Low Birth Weight Rate (<2,500 Grams, Excluding <500 Grams)

Short/Other Names Not applicable

Indicator Description and Calculation

Description

Crude low birth weight rate (<2,500 grams, excluding <500 grams)
 Rate = (Number of live newborns with a birth weight between 500 grams and 2,499 grams, inclusive) ÷ (Number of live newborns with a birth weight greater than or equal to 500 grams) × 100

Calculation: Description

Unit of Analysis: Discharge from acute care facilities

Place of residence

Rate - Rate per 100

None

Not applicable

Calculation: Geographic Assignment

Calculation: Type of Measurement

Calculation: Adjustment Applied

Calculation: Method of Adjustment

Description:

Number of live newborns with a birth weight greater than or equal to 500 grams

Inclusions:

1. ICD-10-CA in-hospital birth codes:

–Any one diagnosis code of Z38.0, Z38.3 or Z38.6 coded in any position

Exclusions:

1. Non-newborn records

2. Cadaveric donors

3. Weight <500 grams

4. Invalid birth weight (Weight = blank, 0000, 0001 or ZZZZ)

5. ICD-10-CA abortion or out-of-hospital birth codes:

–Any one diagnosis code of P96.4, Z38.1, Z38.2, Z38.4, Z38.5, Z38.7 or Z38.8 coded in any diagnosis field

Description:

Number of live newborns with a birth weight between 500 grams and 2,499 grams, inclusive

Inclusions:

Weight <2,500 grams

Exclusions:

Same as denominator (numerator is a subset of the denominator)

Denominator

Numerator

Background, Interpretation and Benchmarks

Low birth weight can add to length of hospital stays and health care costs. Over the fiscal years, these rates have continued to vary among provinces and territories. Given this variability, this indicator can be used to inform government planners, policy-makers, health system

Rationale

| | |
|---|---|
| <p>Interpretation</p> <p>HSP Framework Dimension</p> <p>Areas of Need</p> <p>Targets/Benchmarks</p> <p>References</p> <p>Availability of Data Sources and Results</p> <p>Data Sources</p> | <p>decision-makers, health care providers, patients and the general public. It can help raise awareness and allow Canada to compare its rates with the rates of other countries.</p> <p>Lower rates for this indicator are desirable.</p> <p>Health System Outcomes: Improve health status of Canadians</p> <p>Not applicable</p> <p>Not applicable</p> <p>Not applicable</p> |
| <p>Available Data Years</p> | <p>DAD, HMDB, MED-ÉCHO</p> <p>Type of Year: Fiscal</p> <p>First Available Year: 2001</p> <p>Last Available Year: 2017</p> |
| <p>Geographic Coverage</p> <p>Reporting Level/Disaggregation</p> <p>Result Updates</p> <p>Update Frequency</p> | <p>All provinces/territories</p> <p>National, Province/Territory, Region</p> <p>Every year</p> <p>Web Tool: Quick Stats</p> <p>URL: https://apps.cihi.ca/mstrapp/asp/Main.aspx?Server=apmstrextpri_i&project=Quick%20Stats&uid=pce_pub_en&pwd=&evt=2048001&visualizationMode=0&documentID=029DB170438205AEBCC75B8673CCE822</p> |
| <p>Indicator Results</p> | <p>As of 2015–2016, the national results include non-residents of Canada.</p> |
| <p>Updates</p> <p>Quality Statement</p> | <p>In order to provide a more stable rate estimate, only data from regions with a population of at least 75,000 (from 2001–2002 to 2005–2006) or 50,000 (from 2006–2007 to 2014–2015) was reported. Additional indications for suppression were rate instability and under-reporting. As of 2015–2016, regions are not suppressed based on population; instead, the following suppression methodology is used:</p> |
| <p>Caveats and Limitations</p> | <p>a) Suppression due to privacy: To ensure privacy, a suppression rule is applied to all clinical indicator results. Numerators and/or denominators between 1 and 4 are suppressed.</p> <p>b) Suppression due to unstable results: Unstable indicator results are suppressed. In general, these are indicator results with a denominator between 1 and 49.</p> <p>As of 2006–2007, all provinces and territories submit in the International Statistical Classification of Diseases and Related Health Problems, 10th Revision, Canada (ICD-10-CA) and Canadian Classification of Health Interventions (CCI) coding classification systems; for this reason, Canadian rates are provided only from 2006–2007 onward.</p> |
| <p>Trending Issues</p> | <p>Results are presented for a patient's region of residence, rather than the location of the facility where hospitalization occurred. For Prince Edward Island and the territories, the entire province/territory is included as a region. Health regions in Nova Scotia and Alberta have changed over time. The Postal Code Conversion File (PCCF/PCCF+) was used to assign patients to their region of residence.</p> <p>As of 2015–2016, the national results include non-residents of Canada.</p> <p>Nunavut did not submit data to CIHI in 2002–2003.</p> <p>Data from Region 6 in New Brunswick in 2004–2005 was suppressed due to incomplete data submission. Data from this region was excluded from the provincial rate. Data from Peace Country in Alberta in 2006–2007 and 2008–2009 was also suppressed due to incomplete data submission.</p> |
| <p>Comments</p> | <p>2016–2017 data for Nunavut is incomplete; indicator results have therefore been suppressed.</p> <p>Not applicable</p> |