

Mental Health General Hospital Inpatient Separations as a Percentage (Pan-Canadian) of All General Hospital Inpatient Separations, by Province, Diagnosis Category, Sex and/or Age Group

Name	Mental Health General Hospital Inpatient Separations as a Percentage (Pan-Canadian) of All General Hospital Inpatient Separations, by Province, Diagnosis Category, Sex and/or Age Group
Short/Other Names	Not applicable
Description	<p>Percentage of separations (discharges or deaths) for mental illness/addiction out of all general hospital separations by</p> <ul style="list-style-type: none"> • Jurisdiction (province/territory) • Age group • Sex • Diagnosis category
Interpretation	This indicator provides descriptive information about the proportion of mental illness inpatient separations from general hospitals in Canada stratified by diagnostic, geographic and demographic characteristics.
HSP Framework Dimension	Health System Inputs and Characteristics: Health system resources
Areas of Need	Not applicable
Geographic Coverage	All provinces/territories
Reporting Level /Disaggregation	National, Province/Territory
Indicator Results	https://apps.cihi.ca/mstrapp/asp/Main.aspx?Server=apmstrextpd_i&project=Quick%20Stats&uid=pce_pub_en&pwd=&evt=2048001&visualizationMode=0&documentID=D901470C442A5B26D6CC89B28B21C75F

Identifying Information

Name	Mental Health General Hospital Inpatient Separations as a Percentage (Pan-Canadian) of All General Hospital Inpatient Separations, by Province, Diagnosis Category, Sex and/or Age Group
Short/Other Names	Not applicable
Indicator Description and Calculation	Percentage of separations (discharges or deaths) for mental illness/addiction out of all general hospital separations by
Description	<ul style="list-style-type: none"> • Jurisdiction (province/territory) • Age group • Sex • Diagnosis category
Calculation: Description	Percentage of total separations in general hospitals involving mental illness/addiction = Total separations in general hospitals involving mental illness/addiction divided by total separations in general hospitals involving both mental illness/addiction and non-mental illness /addiction diagnoses, by grouping (such as age, sex, province/territory or diagnosis category) multiplied by 100
Calculation: Geographic Assignment	Place of service
Calculation: Type of Measurement	Percentage or proportion
Calculation: Adjustment Applied	None
Calculation: Method of Adjustment	Not applicable

Description:

Inpatient separations from general hospitals involving mental illness/addiction and non-mental illness/addiction diagnoses

Inclusions:

- Denominator
1. Admission to a general hospital in Canada
 2. Sex recorded as male or female for analyses with breakdown by sex
 3. Valid age recorded for analyses with breakdown by age

Exclusions:

1. Newborns
2. Psychiatric hospital separations

Description:

Inpatient separations from general hospitals involving mental illness/addiction as the primary diagnosis and/or diagnosis category (for data extracted from the Ontario Mental Health Reporting System [OMHRS])

Inclusions:

1. Admission to a general hospital in Canada
2. Sex recorded as male or female for analyses with breakdown by sex
3. Valid age recorded for analyses with breakdown by age
4. Mental illness/addiction as the primary diagnosis and/or diagnosis category (for data extracted from OMHRS)

The following codes were used for inpatient separations involving mental illness/addiction:
For 2011–2012 through 2013–2014

- DSM-IV-TR: 290–319, 607.84, 608.89, 625.0, 625.8 and 780.09
- ICD-9-CM: 290–319 and 648.40–648.44
- ICD-10-CA: F00–F99, G30, R41.0, R41.3 and O99.30

For 2014–2015 through 2015–2016

- DSM-IV-TR: 290–319, 607.84, 608.89, 625.0, 625.8 and 780.09
- ICD-9-CM: 290–319 and 648.40–648.44
- ICD-10-CA: F00–F99, G30 and O99.30

As of 2016–2017

- ICD-10-CA codes: F00–F99, G30 and O99.30
- DSM-IV-TR codes: 290–319, 607.84, 608.89, 625.0, 625.8 and 780.09

DSM-IV diagnostic category

- a = Disorders of childhood/adolescence
- b = Delirium, dementia and amnesic and other cognitive disorders
- c = Mental disorders due to general medical condition
- d = Substance-related disorders
- e = Schizophrenia and other psychotic disorders
- f = Mood disorders
- g = Anxiety disorders
- h = Somatoform disorders
- i = Factitious disorders
- j = Dissociative disorders
- k = Sexual and gender identity disorders
- l = Eating disorders
- m = Sleep disorders
- n = Impulse-control disorders not classified elsewhere
- o = Adjustment disorders
- p = Personality disorders

Numerator

DSM-5 diagnostic category

- a = Neurodevelopmental disorders
- b = Schizophrenia spectrum and other psychotic disorders
- c = Bipolar and related disorders
- d = Depressive disorders
- e = Anxiety disorders
- f = Obsessive-compulsive and related disorders
- g = Trauma- and stressor-related disorders
- h = Dissociative disorders
- i = Somatic symptoms and related disorders
- j = Feeding and eating disorders
- k = Elimination disorders
- l = Sleep-wake disorders
- m = Sexual dysfunctions
- n = Gender dysphoria
- o = Disruptive, impulse-control and conduct disorders
- p = Substance-related and addictive disorders
- q = Neurocognitive disorders
- r = Personality disorders
- s = Paraphilic disorders
- t = Other mental disorders

For information on prior years, please contact the Mental Health and Addictions team at mentalhealth@cihi.ca.

Exclusions:

1. Newborns
2. Psychiatric hospital separations

Background, Interpretation and Benchmarks

Rationale Having a solid picture of the proportions of separations and the trends over time provides health system planners and other decision-makers with information to plan care and evaluate the performance of their health care system.

Interpretation This indicator provides descriptive information about the proportion of mental illness inpatient separations from general hospitals in Canada stratified by diagnostic, geographic and demographic characteristics.

HSP Framework Dimension Health System Inputs and Characteristics: Health system resources

Areas of Need Not applicable

Targets /Benchmarks Not applicable

References Not applicable

Availability of Data Sources and Results

Data Sources DAD, HMDB, HMMDB, OMHRS

Type of Year:
Fiscal

Available Data Years
First Available Year: 2003
Last Available Year: 2018

Geographic Coverage All provinces/territories

Reporting Level/Disaggregation National, Province/Territory

Result Updates

Update Frequency Every year

Indicator or Results **Web Tool:** Quick Stats

URL: https://apps.cihi.ca/mstrapp/asp/Main.aspx?Server=apmstxtrpd_i&project=Quick%20Stats&uid=pce_pub_en&pwd=&evt=2048001&visualizationMode=0&documentID=D901470C442A5B26D6CC89B28B21C75F

2016–2017: For records from the Ontario Mental Health Reporting System (OMHRS) — one of the data sources that populates the Hospital Mental Health Database (HMMDB) — broad mental health category in the HMMDB was assigned based on the DSM-5 diagnosis category instead of the specific diagnosis code.

Update es The assignment of diagnosis codes to diagnosis categories was modified. Several ICD-10-CA and DSM-IV-TR diagnosis codes were regrouped into HMMDB broad mental health categories. These changes decreased the volume of separations in the Anxiety Disorders broad mental health category and increased the volume of separations in the Other Disorders category.

Quality Statement

As this indicator excludes psychiatric hospitals, it should not be construed to represent all mental health and addiction hospital separations in Canada.

Certain cells were suppressed for confidentiality reasons in accordance with CIHI's privacy policy. Totals exclude suppressed values.

Many factors contribute to the observed variations in the analysis of indicators at the provincial/territorial and regional levels. These include, but are not limited to, geography, population health, provincial and regional health service resources, and health services administration. It is very important to consider these factors and the effect they may have on indicator results when conducting comparative analyses. The variation in results between provinces/territories or between health regions suggests systematic differences in

Caveats and Limitations

- Provincial/territorial or regional health policies, practices and resources
- Geography and urban/rural population distribution
- Environmental and socio-economic characteristics
- Types of patients served and acuity of illness
- The range of health services available beyond general and specialty psychiatric hospitals (such as community mental health services)
- Accessibility and awareness of alternative points of entry to the health system

The above list suggests a few of the factors that should be considered when making provincial/territorial and/or regional comparisons.

2011–2012: Additional ICD-10-CA diagnosis codes were added to the extraction criteria for the DAD-HMDB, one of the data sources that populates the HMHDB. This increased the volume of separations.

2014–2015: Two ICD-10-CA diagnosis codes were removed from the extraction criteria for the DAD-HMDB, one of the data sources that populates the HMHDB. This decreased the volume of separations.

Trending

2016–2017: For records from the Ontario Mental Health Reporting System (OMHRS) — one of the data sources that populates the Hospital Mental Health Database (HMHDB) — broad mental health category in the HMHDB was assigned based on the DSM-5 diagnosis category instead of the specific diagnosis code.

The assignment of diagnosis codes to diagnosis categories was modified. Several ICD-10-CA and DSM-IV-TR diagnosis codes were regrouped into HMHDB broad mental health categories. These changes decreased the volume of separations in the Anxiety Disorders broad mental health category and increased the volume of separations in the Other Disorders category.

Comments

Separations by year are based on the date of discharge, not the date of admission. As such, records with invalid discharge dates are not included.

The HMHDB is an event-based rather than a person-based database; individuals may be represented multiple times in the data.